



AMERICAN JOURNAL OF PHARMTECH RESEARCH

Journal home page: <http://www.ajptr.com/>

Evaluation of the toxicity of *Annona muricata* leaf extracts on liver and kidney function and investigation of acute and subacute toxicity in Wistar rats

Aivodji Natacha¹, Ahokpe Melanie¹, Justin Behanzin¹, Sezan Alphonse^{1*}

*1.Laboratory of Biomembranes and Cell Signaling University of AbomeyCalavi, 06 BP 3041
Cotonou, Benin*

ABSTRACT

The present work carried out in the laboratory of Biomembranes and Cell Signaling of the University of Abomey-Calavi in Benin aims to conduct a toxicological study on the ethyl extract of the leaves of *Annona muricata*, with a view to verifying the toxic effects in general, and in particular on the functioning of the kidneys and the liver in the wistar rat. In the present study, the extraction yield of this plant by ethanol was 9.6 ± 1.89 . Then, on the one hand, the acute and subacute toxicities were induced in the rats of the experiment by gavage to our extracts at an interval of 48 hours for 10 days. These rats were distributed in 05 batches. Each batch receiving or not receiving a different dose of the *Annona muricata* extract. On the other hand, the toxic activity on the liver and kidneys was evaluated by daily gavage of 14 rats of 3 rats each using the *Annona muricata* ethyl extract at different doses (0.50 100 and 200 mg / kg). Measurement of biochemical parameters (Urea, Creatinine, ASAT, and ALAT), weight-loss records, carried out during the first experiment showed that there was no significant increase urea, creatinine, and a significant decrease in ALT levels. On the other hand, a significant increase was observed in the ASAT. Following the death of the rat of lot 5 receiving 5000mg / kg, the LD50 was determined (LD 50 = 3750 mg / kg). This LD50 indicates that *Annona muricata* extract is weakly toxic. Sub-chronic administration of the extract confirmed that ASAT was progressively increasing in rats. The decrease in body mass was a good indicator of toxicity. In the second experiment, the measurement of biochemical parameters (Urea, Creatinine, Uric Acid, ASAT, ALAT) revealed an increase in the concentration of urea and creatinine and decreased concentration of uric acid at doses of extracts greater than or equal to 100 mg / kg PC. These results may suggest renal damage that has not been confirmed by the histological study of the organs taken (Liver and kidneys) from experimental rats. However, this study showed an early onset of hepatic involvement at the dose of 100mg / kg, which increased at a dose of 200mg / kg.

Keywords: *Annona muricata*, liver, kidneys, LD50, toxicity, Benin

*Corresponding Author Email: sezco@live.fr

Received 4 October 2017, Accepted 30 October 2017

Please cite this article as: Alphonse S *et al.*, Evaluation of the toxicity of *Annona muricata* leaf extracts on liver and kidney function and investigation of acute and subacute toxicity in Wistar rats. American Journal of PharmTech Research 2018.

INTRODUCTION

Plants, which are vital elements of biological diversity, are used primarily for human well-being (Adjanooun et al., 2000). The African populations are confronted with the emergence of chronic diseases whose treatment and follow-up often constitute a major economic problem; they use traditional medicine. Diabetes, high blood pressure and cancer are among the so-called new diseases for this medicine. Once unknown, these diseases have become increasingly important until they become a real public health problem. In response to the expansion of these high-burden diseases, WHO, in its resolution AFR / RC50 / R3 of 31 August 2000, encouraged African countries to develop regional strategies on traditional medicine; to undertake research on medicinal plants and to promote their optimal use in health care delivery systems. Benin is no exception to the practice of this form of medicine. According to the World Health Organization (WHO), 80 to 85% of Beninese people use traditional medicine in the event of illness. One of the plants used in this context in Benin is *Annona muricata*.

The leaves of *Annona muricata* are used as a diuretic, depurative. They are also used for liver diseases. Their use is also found in the treatment of blisters, edema, edema, hygroma, brucellosis, dropsy and undulating fever. (Adjanooun, E, 2000)

But very often, plants are consumed without taking into consideration their toxicity, their synergism with other substances or even the original medicinal proposal established by medicine.

It is recommended to use a plant only on the advice of a specialist because badly dosed the latter can become highly toxic. It is in this context that the present work is carried out, whose main objective is to study the toxicity caused by the ethanolic extract of the leaves of *Annona muricata* on the liver and the kidney.

Generalities on *Annona muricata*



Figure 1: Leaf and fruit of *Annona muricata*

Biology of *Annona muricata*

Annona muricata also named guanábana, guabano or soursop is a type of tree of the Annonaceae family (Annonaceae), and originally grew in the West Indian islands. Today the tree grows in all the Caribbean and Amazonia. The tree is still green and under normal conditions it reaches 6-10 meters in height and has bright oblong-lanceolate leaves that resemble laurel. The yellow-green flowers with 6 petals appear on the branch opposite the leaves. They spread an Asian fragrance and thus attract flies for pollination. The fruits are armored with large black seeds, the size of a small peanut. The fruit is the size of a big apple, green, soft when it is ripe, it is covered with areolas (small growths often present on the cacti) curved soft. The black seeds contain neurotoxin annonacin. Amazonian Indians use the leaves as a medicine. Its fruits are appreciated by the natives for their refreshing virtues and its leaves are traditionally used to cure many diseases.

Classification

Kingdom: Plantae

Under the reign of: Tracheobionta

Division: Magnoliophyta

Class: Magnoliopsida

Subclass: Magnoliidae

Super-order: Magnolianaes

Order: Magnoliales

Family: Annonaceae

Genre: *Annona*

Species: *muricata*

Acute toxicity

It represents the most spectacular manifestation of the harmfulness of a poison. This fact, which leads us to regard as poisonous any substance which kills violently, results in the rapid death of the individual or of the contaminated populations. Acute toxicity can therefore be defined as that which causes death or very serious physiological disorders after a short period of time following transtectonal, pulmonary or buccal absorption. (Ramade, 1979)

The study of acute toxicity is also qualitative and quantitative of the toxic phenomena that can be encountered after administration of the active substance. This study describes the observed symptoms, including local phenomena. It allows:

- Indication of the maximum dose without toxic effect (DME), ie the highest dose for which no toxic effect is observed compared to the control batch;

□ Notation of the minimum dose for which death occurs in all experimental animals; □ determination of the LD50 with its 95% confidence limits (Ruckebusch, 1981).

Determination of the lethal dose (LD50)

The LD50 is in its simplest form the dose of a compound that causes a mortality of 50% in a population of experimental animals. That is, having received a single administration of a product under well-defined experimental conditions. This determination is based on the assessment of all-or-nothing responses: death or survival of animals. The experimental protocol consists of experimenting on 5 to 6 batches of rats to which increasing doses of the test substance are administered in such a way that the percentage of mortality varies between 0 and 100%. This is because it is impossible to immediately obtain 50% of deaths from a single group. The construction of a curve giving the percentage of mortality as a function of the logarithm of the dose leads to determine the dose that would be the LD50 (Wallace Hayes, 2008)

Different methods for determining the LD50

The LD50 can be determined by two calculation methods, the Dragstedt and Lang Method (1957) and the method of Karber and Behrens (1935). As can be determined by two graphical methods, which are the method of Miller and Tainter, (1944), and the method of Litchfield and Wilcoxon, (1949).

Subacute toxicity

It differs from the previous one (acute toxicity) in that a significant proportion of the population can survive intoxication, although all individuals showed short-term clinical signs on target organs, sometimes reversible and resulting from repeated intake of the toxicant, but at a lower dose than that of acute toxicity (Ramade, 1979).Materiels et Methodes

Extraction

The ethanolic extract of the dried leaves powder of *Annona muricata* was prepared from 134.58 g of powder. The powder was macerated in ethanol (95 °). The maceration with duration 72h during which filtrations were carried out every 24h. The solvent was recovered from the filtrate by evaporation with Rotavapor. The filtrate was placed in an oven at a temperature of 40 ° C. The extract obtained was stored in the freezer until it was used.



Figure 2: a. Powder of leaves of *Annona muricata*, b. Evaporation with ROTA VAPOR, c. Extract obtained

Evaluation of the acute toxicity of the ethanolic extract of *Annona muricata*

For the determination of acute toxicity, the experimental protocol used is that described by Rasekh *et al.*, (2008). The rats were randomly assigned to five batches. One batch is used as a control and the other batches are each treated with a single dose of the ethanolic extract of *Annona muricata*. At the beginning of the experiment the rats had an average weight of 200 ± 30 .

o Administration of extracts

The gavages are done every 48 hours and this for 10 days. The doses administered were 0, 100, 1000, 2500, 5000 milligrams of extract per kilogram of rat body weight respectively to the 5 batches.

o Observations

After administration of the extract, the rats are monitored continuously each day. We took notes on apparent signs of toxicity. The purpose of this monitoring was to report changes in diet and water consumption, or behavioral or clinical signs of toxicity.

o Expression of results

The LD50 expressed in mg / kg of body weight is determined by the calculation method of Dragstedt *et lang*, (1957).

This method is based on the following assumption:

Any animal that has survived a dose administered to it will survive at any lower dose. Any animal which has succumbed to a dose administered to it will succumb to any higher dose than the latter. Thus, the percentage of mortality (M [%]) can be calculated for each dose by cumulating all observed deaths at lower doses and all survivors observed at higher doses.

$$M(\%) = \frac{(\text{Number of cumulative deaths})}{(\text{Number of living cumulated} + \text{number of cumulative deaths})} \times 100$$

The LD50 is calculated by interpolation:

$$DL50 = \frac{50(X2-X1) + (X1Y2-X2Y1)}{Y2-Y1}$$

X2: Upper dose framing the LD50; X1: Lower dose framing the LD50; Y2: Percentage of mortality corresponding to X2; Y1: Percentage of mortality corresponding to X1

Evaluation of the subacute toxicity of the ethanolic extracts of *Annona muricata*

For the evaluation of subacute toxicity, four batches of rats were used. The treatments were administered orally once every 48 hours to the rats for 10 days. The first group of animals serving as controls received water, while the other batches received the extracts in respective doses. Lot 2 received 100 mg / kg, lot 3 received 1000 mg / kg, lot 4 received 2500 mg / kg of the extract. Animals are given free tap water and food throughout the test period; and do not receive any other drug treatment.

o Method of observation and examination

Several methods have been used to produce the toxicity texts. These include observation of weight evolution and biochemical tests.

Chronology of weight change

As weight evolution is a good indicator of toxicity, we have followed the weight evolution of rats during our work. Thus, at each 48 hour, the rats of each batch were weighed and their body mass measured in order to see the effect of the extract on the weight evolution.

Blood Collection

Rats are taken before each gavage. The principle of the sample is as follows: the rats are anesthetized with diethyl ether and using a hematocrit tube. the samples are taken from the vein in the angle of the eye. Two tubes were used in EDTA tubes and dry tubes. These tubes were centrifuged at 3000t / 15min and the serum obtained was stored at -15 ° C until use (Ben-Romdhane et al., 2003).

Biochemical examinations

The biochemical analyzes were carried out in the Laboratory of Biochemistry of the Faculty of Health Sciences using a spectrophotometer: Kenza Max distributed by Biolabo. The assay parameters are ALAT (Alanine amino transferase), ASAT (Aspartate amino transferase), CREA (Creatine) and Urea.

Procedure used for the measurement of biochemical parameters

Creatinine

Table 1: Mode of measurement for creatinine

Measure in a vessel of 1cm optical path.	White (optional)	Standard	Sample
Work reagent (R1+R2)	1ml	1ml	1ml
Demineralized Water	100µl		
Standard		100µl	
Specimen			100µl
Mix thoroughly after 30 seconds, record absorbance A1 at 490nm (480-510) against reagent blank or distilled water. Read the absorbance.			

UREA

More than 90% of the urea is eliminated by the kidneys in the urine. Measurement of plasma or serum concentration of urea is often considered an indicator of renal function. However, certain non-renal factors also influence the concentration of urea: urea is increased, among other things, in cases of accelerated protein catabolism (burns, trauma, myocardial infarction ...). The urea level is lowered to the terminal stage of severe hepatic insufficiency and is then accompanied by an increase in ammonia. The urea level is generally studied in conjunction with the creatinine ratio (urea / creatine ratio) to refine the diagnosis of post-renal or pre-renal azotemia. (Pirro, 2014)

Table 2: Mode of measurement of urea

Measure in a thermostatically controlled tank (30 ° C or 37 ° C)	Standard	Sample
Reagent	1ml	1ml
Standard	5µl	
Specimen		5µl
Mix. Read the concentrations at 340nm. 1st reading A1 in 30 seconds. 2nd reading A2 at 90 seconds.		

ALAT**Table 3: Mode Used for ALT Measurement**

Introduce into a reading tank of 1cm optical path:	
Reagents	1 ml
Allow the temperature to equilibrate to 37 ° C and then add:	
Samples	100 µl
Mix. After 1 minute, read the absorbance at 340 nm every minute for 3 minutes.	
Calculate the average absorbance variation per minute (Abs / min variation)	

ASAT**Table 4: Mode Used for ASAT Measurement**

Introduce into a reading tank of 1cm optical path:	
Reagents	1 ml
Allow the temperature to equilibrate to 37 ° C then add:	
Samples	100 µl
Mix. After 1 minute, read the absorbance at 340 nm every minute for 3 minutes.	

Calculate the average absorbance variation per minute (Abs / min variation)

Evaluation of the impact of the ethanolic extract of *Annona muricata* on the functioning of the liver and kidneys at the usual doses.

The experimental protocol is carried out on male adult rats of the "Wistar" type of about 200 g, raised at the laboratory of biomembranes labels and cell signaling at the University of Abomey-Calavi. The animals have been acclimatized from birth to laboratory conditions. They were fed with protein-rich granules, lipids and had available, tap water seamlessly in small jars and have free access to food and water. The breeding is carried out in a room illuminated 12 hours a day, and whose temperature is ambient. These animals will be subjected to sub-chronic gavage (14 days). (OECD. 2009).

The rats are weighed and randomly assigned to 04 batches of 03 rats (No. 1, No. 2 and No. 3), and then the batches are assigned respectively to doses per kg of body weight per day: 0 mg / kg; 50 mg / kg; 100 mg / kg; 200 mg / kg; and the volume to be administered is fixed at V = 01 ml.

Batch 1 is the control batch and receives by oral administration of the distilled water in place of the ethyl extract (0 mg / Kg) of *Annona muricata*, for 14 days.

The other batches (2-4) are treated by oral administration, at the respective doses of 50 mg / kg; 100 mg / kg; 200 mg / kg; of body weight and per day of the *Annona muricata* ethyl extract, for 14 days;

o Blood sampling

The biological material consisted of a blood sample taken from wistar strain albino rats during the various experiments. The blood was collected by puncture at the retro orbital sinus using a micro tube during the following days (j0, j7, j14). The blood is taken from a dry tube (for biochemical examinations). Samples for biochemical examinations were centrifuged at 3000 rpm for 5 min.

o Determination of biochemical parameters

Uric acid

Table 5: Mode used for measurement of uric acid

Measure in well identified test tubes.	White	Standard	Sample
Work reagent	1ml	1ml	1ml
Specimen (Rq1)	-	-	25µl
Standard	-	25µl	-
Demineralized Water	25µl	-	-
Mix, let stand 5 minutes at 25 ° C. Read absorbances at 520nm (490-530) against reagent blank. Staining is stable for 30 minutes.			

For the other parameters: ALAT (Alanine amino transferase), ASAT (Aspartate amino transferase), CREA (Creatine) and Urea; the method used is the same as that described above in the first experiment.

o Histological study of the organs taken (kidneys and liver)

After the last sampling on the 14th day, all surviving animals are sacrificed and dissected in order to observe any signs of tissue toxicity (kidneys and liver).

Procedure of the histological study

Preparation of cassettes

The organs resulting from the dissection were cut into small pieces and then placed in cassettes for fixation. Sharp instruments are used in order not to crush the tissues and thus avoid the formation of artifacts (the scalpel). The sampling is done as quickly as possible.

Fastening

Its purpose is to preserve structures and harden parts. Indeed, the tissue collection causes them to die: the cells discharge their enzymes, which causes a self-regulation of the tissue. Moreover, in the ambient air, the samples can be contaminated by bacteria, which leads to putrefaction of the tissues. It must be done immediately after sampling, by immersing the material in a large volume of fixing liquid. The most frequently used fixing fluids are 4% formalin or Bouin liquid (picric acid + formol + acetic acid water).

The choice of fixer was made on 10% buffered formalin for morphological studies because it is easy to prepare and stable. Moreover, this fixator penetrates the tissues well. The preservation can be carried out in this solution for a relatively long time without the tissues deteriorating. The duration of the fixation is on average of 3 days for the cuts of small sizes.

Interest of fixation:

Fixation is the primary step even in the histological study because it has the advantage: immobilization of tissue / cellular constituents; prevents cellular autolysis; prevents post-mortem bacterial putrefaction; allows the histological technique and subsequent colorations

Its principle is based on the fact that it reacts with amino groups of proteins. The fixation time is variable and the amount of fixer used must be at least ten times greater than the volume of tissue to be fixed: a few hours are sufficient to fix the small fragments. It has the disadvantage of causing a phenomenon of cellular auto-fluorescence which can hamper observations under the fluorescence photon microscope. It also rapidly degrades the nucleic acids.

Role of the fixer:

The role of the fixer rests on actions such as:precipitation, polymerization, coagulation of proteins;establishment covalent bonds;kills the cells;blocking of enzymatic reactions.

Inclusion or circulation

The inclusion allows fine and regular cuts. In order for the light to pass through the tissue to be examined, it must be very thin. Now the tissues are soft, so they must be given a solid consistency. This is the principle of inclusion: paraffin inclusion involves infiltrating and coating the tissues to be examined with paraffin. Inclusion is preceded by two essential steps. First of all,

dehydration: the tissues are passed into alcohol baths of increasing degree (70 °, 80 °, 90 °, 95 °, 99 °, and finally 100 °). The interest of dehydration is to eliminate the fixative.We therefore used as a dehydrating agent ethanol because it has a great miscibility with water and allows a good hardening of the tissues. Dehydration was progressive, by passing the tissue pieces into alcohol baths of increasing concentrations, to avoid distortion of the tissue structures. We used seven ethanol baths of increasing concentration. The rooms stayed for an hour in each of these baths.

The alcohol (ethanol) is then replaced by a solvent which is miscible with paraffin: it is either xylene or toluene (hydrocarbons). These substances eliminate ethanol. As they are infiltrated by the solvent, the tissues tend to become lighter: this stage is sometimes called clarification or clarification.

The choice of our solvent is based on xylene, we have carried out three (03) baths there. Our manipulations were done under a chemical hood. As soon as the first xylene bath begins to become cloudy (which translates to incomplete dehydration of the room), we immediately remove the tissue and plunge it into a new bath of absolute ethanol. The rooms have stayed about an hour in each of the baths.

The impregnation, the fabric is placed in molten paraffin (raised to 56/58 ° C); the heat causes the solvent to evaporate (and its dissolution in the paraffin): the spaces thus released are filled with paraffin. We have here used three paraffin baths, the rooms have stayed about an hour in each of the baths.

Embedding

The paraffin is placed in small molds at ambient temperature, which causes it to harden and thus to stiffen the tissue fragments taken. The demoulding is then carried out: tissue fragments included in a paraffin block are obtained. We have deposited and oriented tissues in metal molds to facilitate further study. The molten paraffin is poured into the molds which support the blocks to be formed.

Finally the mold block assembly is set to cool in a refrigerator. Once the block of paraffin has solidified, it is separated from its mold and is then ready to be cut to the microtome.

Microtomy

We then isolate cuts in the paraffin block. We use a microtome, which advances the block on a razor: the block advances about 2 to 3 μm each time (in classic MO, the cuts measure about 5 μm).

The whole of the slices will form a ribbon in which there are serial sections for tissue sampling.

To do this we have inserted the block into the object clamp and then immobilized by means of the locking screw and oriented with the adjustment screws, making sure that the bottom and top faces of the block are parallel to the cutting edge of the razor. After checking the parallelism between the block and the razor, the motor wheel of the microtome is turned by means of the crank.

□ Spreading and sticking of sections on glass slides

We heat the paraffin on a hot plate and so the paraffin sticks to the blade. We cover blades clean and engraved with glycerine water. Successful ribbon segments are then selected using lancets and placed on slides. The whole is deposited on the heating plate and the ribbons are allowed to stretch. With two lancets we pull on the edges of the tissues to eliminate the persistent folds; the excess liquid is absorbed with a blotting paper and the ribbons are allowed to adhere to the slides. These are then placed in an oven at 40 ° C. for one hour after drying; finally they are allowed to cool for at least 15 minutes. The preparation is then ready for coloring.

□ Staining of blades

Made on slides, the coloring accentuates the contrasts to better recognize the different elements of the preparation. The tissues of the organism are not spontaneously colored, which makes the observations difficult. The dyes used in histology are more or less selective; most are acidic or basic components in an aqueous medium which form salts with the ionized radicals of the tissues. Acidic components are used for basophilic tissue areas, and basic components are used for acidophilic tissue areas. The most commonly used staining is HES: hematein / eosin / saffron. Hematein is a rather basic substance, which colors the nuclei in violet thus coloring the nucleic acids. Eosin is a rather acidic substance, which rather colored the cytoplasm (in pink) thus colored the proteins. Finally, the saffron colors the collagen fibers in yellow.

However in order for a stain to be used, paraffin must be removed. Dewaxing is then performed, which consists of passing the slides in toluene or xylene baths in order to dissolve the paraffin. The mixture is then rehydrated: the alcohol is mixed with water and toluene, and the slides are passed through alcohol baths of decreasing degree (from 100 ° to 70 ° or even 50/40 °). You can also use

histochemical staining, called PAS (Periodic Acid Schiff), which highlights glycogen, glycoproteins and just about anything that contains sugars.

The coloring chosen in the case of our study is that with haematein eosin. The haematein staining lasted 15 min, the running water bath → 5 min, the acidic alcohol bath → 2 to 3 dives, the running water bath → 5 min, the ammonia water bath → 30sec to 1 min, the running water bath → 5 min, the eosin stain: 15 sec → 2 min, the running water bath → 5 min.

Blade assembly

The assembly is carried out by affixing the object slides to the histological slide using Eukitt glue.

Reading

Microscopic observation is here coupled with photomicrography. Our observations were made on a light microscope equipped with a camera. The images were transferred to a processing software for color to be used, the paraffin must be removed. Dewaxing is therefore carried out, which consists in passing the slides in baths of toluene or of xylene in order to dissolve the paraffin. The alcohol is then mixed with water and toluene, the slides are passed into alcohol baths of decreasing degree (from 100 ° to 70 °, or even 50/40 °). One can also resort to a histochemical staining, called PAS (Periodic Acid Schiff), which highlights glycogen, glycoproteins and just about anything that contains sugars.

The coloration chosen in the case of our study is that to hematein eosin. The hematein staining lasted 15 minutes, the bath of running water → 5 minutes, the acid alcohol bath → 2 to 3 dives, the water bath → 5 minutes, the ammonia water bath → 30 seconds to 1 mn, running water bath → 5 mn, staining with eosin: 15 sec → 2 mn, running water bath → 5 mn.

Assembly of the blades

The assembly is made by affixing the object slides on the histological slide with Eukitt glue.

Playback

Microscopic observation is here coupled with photomicrography. Our observations were made on a photonic microscope equipped with a camera. The images were transferred to image processing software (Adobe Photoshop Image) using a digitizer that transforms the analog image into a digital image. The cuts are observed at the appropriate magnifications; the images are captured according to the desired objective and are transferred in JPG format.

RESULTS AND DISCUSSION

Evaluation of extraction techniques

The ethanolic extract of *Annona muricata* was obtained from a fine powder of the leaves of the

corossolier. The method consisted in carrying out a maceration of the *Annona muricata* powder in ethanol for 72 hours and every 24 hours was carried out the evaporation of the solvent to the rotavapor. The results of the present study indicate that from 134.58 g of *Annona muricata* powder and 1 liter of ethanol we obtained an ethanol extract, considered as the dark green colored extract which may contain chlorophyll, flavonoids, polyphenols and other compounds. In order to use the extract for pharmacological and toxicological tests, the conservation of the bioactive state of the extracted molecules seems important. Complete depletion of the solvent is necessary. The presence of traces of ethanol in the extract can cause undesirable side effects. Therefore, the positive or curative effect of the pharmacological substance can be masked by the action of the solvent. The yield was calculated on the total weight of the powder of the *Annona muricata* powder and that of the ethanol extract gave a yield of $9.60 \pm 1.89\%$. This result is close to that found by AÏVODJI N., (2014) who used the same extraction protocol for the leaves of the same plant.

Study of Acute Toxicity

o Determination of LD 50

Before attempting to determine the LD50, a preliminary search for LD0: the highest dose at which all experiencing animals survive, and LD100: the lowest dose at which all animals die, is required. These two doses delimit the area in which the final test is to be performed. In our work the test is carried out according to the method described by Rasekh *et al.*, (2008), on male Albino Wistar white rats. Five (05) doses, including DL0 and DL100, were retained (Table 6), while control rats (0mg / kg) remained healthy.

Table 6: Doses injected and percentage conversion of the number of dead rats in each batch.

Lot	Number of rats (n)	Doses (mg/kg)	Number of dead rats	Percentage (%)
1	1	0	0	0
2	1	100	0	0
3	1	1000	0	0
4	1	2500	0	0
5	1	5000	1	100

After calculating the percentages of mortality in each batch, we can then determine the LD50

o Calculation method or method of Dragestedt and Lang. After applying the relationship:

$$M(\%) = \frac{(\text{Number of cumulative deaths})}{(\text{Number of living cumulated} + \text{number of cumulative deaths})} \times 100$$

The LD50 is calculated by interpolation:

$$LD50 = \frac{50(X2 - X1) + (X1Y2 - X2Y1)}{Y2 - Y1}$$

X2: Upper dose framing the LD50; X1: Lower dose framing the LD50; Y2: Percentage of mortality corresponding to X2; Y1: Percentage of mortality corresponding to X1. This method revealed an LD50 equal to: 3750 mg / kg.

o Characterization of the form of toxicity of the ethanolic extract of *Annona muricata*

The highest dose killing all animals or 100% lethal dose (LD100) is 5000 mg / kg and the maximum tolerated dose (DMT), the maximum dose that does not kill any animal when the extract is administered, is 2,500 mg / kg /. In the numerical application of the formula of Dragestedt and Lang (1957), the calculation of the LD 50 of the extract gives the value of 3750 mg / kg. According to the Diezi (1989) toxicity scale, this LD50 value (3750 mg / kg) indicates that the ethanolic extract of *Annona muricata* leaves, administered orally (VO), is weakly toxic in rats under the conditions of this study.

Table 7: Toxicity class, according to the Diezi toxicity scale (1989)

Index or class of toxicity	Common term used	Toxicological parameter (LD50)
1	Very toxic substance	LD50 < 5mg/kg
2	Toxic substance	5mg/kg < LD50 < 500mg/kg
3	Low toxic substance	500mg/kg < LD50 < 5000mg/kg
4	Non-toxic substance	LD50 > 500mg/kg

In this study, the LD50 of the ethanolic extract of *Annona muricata* calculator is included between 500 mg / kg and 5000 mg / kg (500 mg / kg < 3750 mg / kg < 5000 mg / kg). The ethanolic extract of *Annona muricata* is considered to be low toxicity in rats by oral route according to the classification of Diezi, (1989) and therefore this plant deserves to be used with caution in human

Study of Acute Toxicity

After administering the oral ethanol extract of *Annona muricata* to male rats, there is a steady increase in signs of poisoning and mortality in relation to the dose injected. The signs are summarized in the table.

Table 8: Sign of poisoning observed at each dose

Doses (mg/kg)	Rats Lots	M/T	Toxicity symptoms
0	Lot1: witness	0/1	Nothing
100	Lot 2: a blue bar	0/1	No visible signs toxicity
1000	Lot 3: a red bar	0/1	Hair straightening, drowsiness, loss of appetite, weakness
2500	Lot 4: a green bar	0/1	Hair straightening, drowsiness, loss of appetite, weakness
5000	Lot 5: 2 red bars	1/1	Hair straightening, drowsiness, loss of appetite, weakness, rapid heartbeat, labored breathing, body tremor, loss of balance, convulsion, total paralysis, coma, death

The first dead rat was recorded at a dose of 5000 mg / kg after 4 injections following severe signs of intoxication. After the second injection, the rat showed almost cut breathing, loss of balance, muscle contractions, partial paralysis (Fig 3-c), coma and total paralysis, he died on the 6th day. All the other rats survived. However, signs of toxicity have been noted. Namely hair straightening, drowsiness, motor weakness and loss of appetite. The dose that kills the entire rat population is 5000mg / kg, while the 2500mg / kg dose is tolerated. The signs of toxicity increased proportionally with the doses administered. Thus we observed more severe signs of toxicity in rats given 2500 mg / kg. Gradually we noticed that the movement of rats as well as their appetite decreased and that their breathing became difficult. At the 1000mg / kg dose, the percentage of death is nil however we noted a motor weakness which limited the rat in its movement. As for the 100 mg / kg dose, it did not show signs of tangible toxicity, however, this dose could be useful in the study of long-term toxicity.



Figure 3: Acute signs of acute toxicity in rats. (a) Straightening of the hairs; (b) rats are drowsy and tired; (c) Partial paralysis at dose 5000mg / kg

Subacute toxicity study

o General signs

No deaths were recorded during the experiment. However, from a behavioral point of view, the rats are calm, showing a decrease in liveliness and appetite compared to the control rat; their sleep time has become more considerable.



o Chronology of weight change

The change in animal body mass during the subacute toxicity experiment with the ethanol extract of *Annona muricata* showed that there was a decrease in weight throughout the 2 weeks, compared to witness (Figure. 7).




Figure 5: Changes in body weight of rats as a function of time under subacute toxicity conditions by the ethanolic extract of *Annona muricata*.

The percentage decrease was minimal during the first six days of the experiment. This percentage was 4.08% for lot 2 receiving 100 mg / kg; there was no decrease in lot 3; at Lot 4 a decrease of 2.43% was observed. These decreases are greater on the eleventh day of the experiment and were 18.36% for lot 2; 8.10% for lot 3 and 14.63% for lot 4. (Tab 10).

Table 10: Percentage decrease in body weight of rats during the period of subacute treatment with ethanolic extract of *Annona muricata*.

Days	Lot2	Lot 3	Lot4
6	4,08%	0%	2,43%
11	18,36%	8,1%	14,63%

The change in body weight is used as an indicator of the adverse effects of chemical compounds (Hilaly et al., 2004). Weight loss is correlated with the physiological state of the animal. These decreases in body weight in animals may be due to lower food consumption, we can also relate this decrease in weight to the possibility of dose / absorption interactions and the decrease in the amount of food intake. The decrease in body weight during the treatment period suggests that subchronic and oral administration of the ethanolic extract of *Annona muricata* has effects on the growth of Albino Wistar rats. To make this hypothesis more precise, other analyzes deserve to be carried out.

o Biochemical data

Several toxic plant compounds accumulate in the liver where they are detoxified. The liver is the first target of toxicity and the first organ exposed to all that is absorbed in the small intestine; it metabolizes foreign substances to compounds that may be hepatotoxic. (P. Hanover, 1963) The liver works in combination with the kidneys to remove toxic substances from the blood (Tulsawani, 2010). A study of renal and hepatic function may therefore be useful for assessing the toxic effects of medicinal plants. These tests mainly include the determination of ASAT, ALAT, CREA and others (Tilkian, 1979) and any necrosis of liver cells leads to a significant increase in ASAT, ALT, and serum enzymes (Adeneye et al., 2006).). Serum studies in rats treated with the ethanolic extract of *Annona muricata* show a significant increase in the ASAT parameter in rats and a significant decrease in ALT in rats. The other parameters: CREA, Urea do not show any significant changes.




Figure 6: Changes in serum creatinine levels of rats as a function of time under subacute toxicity conditions by the ethanolic extract of *Annona muricata*.




Figure 7: Changes in serum urea level of rats as a function of time under the conditions of subacute toxicity by the ethanolic *Annona muricata* extract.




Figure 8: Variations in serum ALT levels of rats as a function of time under subacute toxicity conditions by the ethanolic extract of *Annona muricata*.




Figure 9: Changes in serum ASAT levels of rats as a function of time under subacute toxicity conditions by the ethanolic extract of *Annona muricata*.

ASAT is an enzyme considered to be a good indicator of liver function (Hilaly et al., 2004) and a bio marker to predict possible toxicity. In general the elevation of transaminases in the blood is an

index of damage of the parenchymal cells of the liver. In addition, ASAT found in serum is of mitochondrial and cytoplasmic origin, and any increase can be taken as a first sign of cell damage that leads to the flow of the enzyme into the serum. Thus, the significant increase in ASAT strongly suggests that subchronic administration of *Annona muricata* alters the hepatocytes and hence the metabolism of the rat. ASAT is an enzyme with a high concentration in hepatocytes, but it is also widely distributed in the brain, kidneys, skeletal muscles and myocardium. The increase of this enzyme is an indication of the likely toxicity of these organs and requires further analysis. Evaluation of the impact of the ethanolic extract of *Annona muricata* on the functioning of the liver and kidneys at the usual doses. The results of the analysis of the variance showed a very highly significant difference between the different doses of the extract with respect to the amount of creatinine in the blood (Prob. <0.001). Similarly, the time on this parameter is very highly significant (Prob. <0.001). However, there was no significant difference in the interaction between doses of the extract versus time on this parameter (Prob. > 0.05).




Figure 10: Variation of creatinine by doses of *A. muricata* ethyl extract

The Student-Newman-Keuls test (SNK) made it possible to form homogeneous groups of extract with respect to creatinine. A high creatinine value was observed with the use of the Ext100 mg / Kg extract followed by the Ext200 mg / Kg. On the other hand, the control treatment gave the lowest creatinine value.

The SNK test showed that the highest creatinine value was observed on day 14, followed by the 7th day after administration regardless of dose of the extract.

UREA

The results of analysis of the variance showed a very highly significant difference between the different doses of the extract with respect to the amount of urea in the blood (Prob. <0.001). Similarly, the effect of time on this parameter is very highly significant (Prob. <0.001). However, there was no significant difference in the interaction between doses of the extract and time-dependence on this parameter (Prob. > 0.05).




Figure 11: Variation of the urea according to the doses of the ethyl A extract. muricata

The SNK test made it possible to form homogeneous groups of extract with respect to the amount of urea in the blood. A high value of urea is observed with the use of the Extra200 mg / Kg extract followed by the Ext100 mg / Kg. On the other hand, the control treatment gave the lowest value of the urea in the blood. The SNK test showed that the highest value of urea was observed on the 14th day and 7th day after administration regardless of the dose of the extract.

□ Uric Acid

The results of analysis of the variance showed a significant difference between the different doses of the extract with respect to the amount of uric acid in the blood (Prob <0.05). Similarly, the time factor on this parameter is significant (Prob <0.05). On the other hand, there is a significant difference in the interaction between the doses of the extract and the time dependence on this parameter (Prob <0.05).




Figure 12: Variation of the ASAT according to the doses of the ethyl A extract. muricata

The SNK test showed that the highest value of ASAT on day zero and on the 7th day after administration whatever the dose of the extract.

ALT

The variance analysis results showed no significant difference between the different doses of the extract with respect to ALT (Prob > 0.05). The time on this parameter is very highly significant (Prob. <0.001). On the other hand, there is a very highly significant difference in the interaction between the doses of the extract and as a function of time on this parameter (Prob. <0.001).




Figure 13: Variation of the ASAT according to the doses of the ethyl A extract. muricata

The SNK test showed that the highest value of ASAT on day zero and on the 7th day after administration whatever the dose of the extract.

□ ALT

The variance analysis results showed no significant difference between the different doses of the extract with respect to ALT (Prob > 0.05). The time on this parameter is very highly significant (Prob. <0.001). On the other hand, there is a very highly significant difference in the interaction between the doses of the extract and as a function of time on this parameter (Prob. <0.001).




Figure 14: Variation of the ALT according to the doses of the ethyl A extract. muricata

The SNK test showed that the greatest value of ALT on day zero and on the 7th day after administration whatever the dose of the extract. The SNK test revealed that the highest ALT values were observed in the control rats on day zero irrespective of the dose of the extract. On the other hand, the lowest values of the ALT are observed with the doses of the extra extract 100mg / Kg, ext 200mg / Kg and the control on the 14th day.




Figure 15: Hepatic control rat lobule showing its typical architecture with hepatocyte trabeculae separated by vascular spaces and arranged in radius around the centrilobular vein (VC) (HES × 40)




Figure 16: Renal cortex of control rats showing a renal glomerulus with flocculi and urinary tract and urinary tubules (HES \times 40)




Figure 17: rat hepatic lobule of lot 1, which received 50 mg / kg of PC of the *Annona muricata* ethyl extract: the hepatic architecture is generally conserved (HES \times 40)




Figure 18: Batch 1 rat kidney cortex receiving 50 mg / kg PC of *Annona muricata* ethyl extract: no significant structural changes (HES \times 40)




Figure 19: Rat hepatic lobule from lot 2 which received 100 mg / kg PC of *Annona muricata* ethyl extract: although hepatic architecture remains recognizable, hepatocytes are atrophied and their nuclei are pycnotic (HES \times 40)




Figure 20: Batch 2 renal rat cortex receiving 100 mg / kg PC of *Annona muricata* ethyl extract: almost normal histological appearance (HES \times 40)




Figure 21: Rat hepatic lobule of lot 3 which received 200 mg / kg of PC of the *Annona muricata* ethyl extract: in addition to pycnosis, it should be noted that some hepatocytes are in hydropic degeneration (HES \times 40)




Figure 22: Rat Renal Cortex of Lot 3 receiving 200mg / kg PC of *Annona muricata* ethyl extract: there are no morphologically identifiable abnormalities (HES \times 40)

The study of the biochemical parameters showed that the extract at various doses had no effect on the liver cells since this had no influence on the biochemical parameters (ASAT and ALAT) during the 14 days in the blood (these results were consistent with that of Bridgemohan P., 2013). On the other hand, the histological study showed that from the dose 100 mg / kg PC of the extract, hepatic involvement was observed. The latter is accentuated in a hepatic pycnosis of the nuclei at the dose of 200 mg / kg PC, with an erasure of the cellular limits in the hepatocytes without modification of the architecture. These results confirm those of our first experiment, in which hepatic involvement was suspected in acute toxicity. The variance analysis results showed a very

highly significant difference between the different doses of the extract with respect to the amount of creatinine and urea in the blood. A high value of creatinine and urea was observed with the use of extract Ext100 mg / Kg followed by Ext200 mg / Kg in blood compared to the time of administration of the extract. However, there is no change in values or toxic effect on these parameters with the use of the extract Ext50mg / Kg. In terms of uric acid variance analysis results, there was a reduction in the time-dependent rate of administration of the Ext50mg / Kg extract, which would explain the results of Bridgemohan P. (2013). Therefore, at a dose greater than or equal to 100mg / Kg PC, there would be damage to the renal tissues. However, the histological study showed that at concentrations of 50,100 and 200 there is no negative effect on the kidneys. The animals in general have supported the usual doses allowed. No deaths have been recorded. All animals survived at the end of the 14 days of observation, which did not determine the LD50. This would imply that the LD50 is greater than 200mg / kg bw, and confirm our previous results which set the LD50 at 3750mg / kg PC extract.

CONCLUSION

Our study is based on the species *Annona muricata*, a plant widely used in Benin for its therapeutic properties in the treatment of hypertension, diarrhea, fever, flu and spasms. The continued use of this plant can lead to side effects that can extend to the death of the individual. The risk of using this plant would be closely correlated with the doses administered. In this work, the toxicity of *Annona murica* has been studied at two levels: acute toxicity and subacute toxicity of the plant's ethanol extract. Information from Acute toxicity, subacute toxicity and even routine doses (50,100 and 200) in the Albino Wistar rats suggest that *Annona muricata* the category of low-toxic plants orally. The ethanolic extract of *Annona muricata* may cause symptoms of toxicity that are dose-dependent, ranging from mild sleepiness, loss of appetite, breathing problems, and accelerated heart rhythm, leading to seizures , tremors, comas and paralysis of the legs and even total paralysis leading to death. This toxicity also results in decreased weight, and liver damage. The results of our work give us an idea of the toxicity of this plant. However, further studies are desirable.

REFERENCES

1. Adjanooun E., de Souza S., et Sinsin B. 2000. La biodiversité face au développement des industries pharmaceutiques africaines. In : Réseau des « espèces ligneuses médicinales », EyogMatig O (eds). Compte rendu de la première réunion du réseau tenue 15-17 décembre 1999 à la station IITA Cotonou, Bénin, 88-103

2. Ramade, F., 1979. Ecotoxicologie, Ed Masson, Paris, pp. 5.
3. Ruckebusch, Y., 1981. Physiologie, pharmacologie, thérapeutique animale, Ed. Maloine, Paris, p. 611.
4. Wallace Hayes, A., 2008. Principle and methods of toxicology. Ed Tayler& Francis, New York, p. 1134.
5. Ben-Romdhane, S., Romdane, M.N., Feki, M., Sanhagi, H., Kaabachi, N., M'Bazaa, A., 2003. Valeurs usuelles des principaux constituants biochimiques sériques du dromadaire (*Camelusdromedarius*). Méd. Vét154(11), 695-702.
6. Rasekh, H.R, Khoshnood-Mansourkhani, M.J., Kamalinejad, M., 2001
7. Dragsted, A., Lang, B., 1957. Etude de la toxicité par administration unique d'un nouveau médicament. Annales pharmaceutiques Française, p.11.
8. Diezi, J., 1989. Toxicologie : principes de base et répercussions cliniques. Ed Slatkine, Genève, pp. 33-44.
9. Hilaly et al., 2004. Hypolipidemic effects of Teucriumpoliumin rats. Fitoterapia72, 937-939.
10. Tulsawani, R., 2010. Ninety day repeated gavage administration of Hippophaeerhamnoides extract in rats. Food and ChemicalToxicology48, 2483– 2489.
11. Tilkian, S.M., 1979. Clinical Implications of Laboratory Tests. The C.V. Mosby Company, Missouri, pp. 11–17.
12. Adeneye, A.A., Ajagbonna, O.P., Adeleke, T.I., Bello, S.O., 2006. Preliminary toxicity and phytochemical studies of the stem bark aqueous extract of Musangacecropioidesin rats. Journal of Ethnopharmacology105, 374–379.
13. Karber, C., Behrens, B., 1935. WiesindReihenversuche fur biologischeAuswertungen am ZweckmässigstenAnzuordnen. Arch. Exp. Path. Pharm177, 379-388.
14. Litchfield, J.T., Wilcoxon, F.A., 1949. A simplified method of evaluating dose-effect experiments. J. Pharmacol. Exp. Ther96, 99-113.
15. Miller, L.C., Tainter, M.L., 1944. Estimation of ED50 and its error by means of logarithmic. Probitpaper. Proc Soc Exp Viol Med 57, 261-4.
16. OCDE. (2009). PROJET DE LIGNE DIRECTRICE DE L'OCDE POUR LES ESSAIS DE PRODUITS CHIMIQUES (Section 4 : Effets sur la santé ; Essai N°453 Études combinées de toxicité chronique et de cancérogenèse, adoptée le 08 Septembre 2009).
17. Pirro, 2014. Reins et voies urinaires-appareil génital masculin : anatomie du rein et vascularisation

18. P. HANOVER (1963 - 1964), METHODES D'ANALYSES utilisées au Laboratoire des Glucides c.s.T. BONDY
19. Natacha A. 2014, Action d'extraits éthyliques de *Annona muricata* dans le traitement traditionnel du cancer de foie. Screening phytochimique et recherche de l'activité antiproliférative chez le rat wistar. Mémoire de fin de formation en Master de Physiologie et Pharmacologie Cellulaires.
20. Bridgemohan P., Bridgemohan R.S.H., 2013; Evaluation of decoction extracts of two types of soursop (*Annona nuriata*) for annonaceous acetogenin properties.

AJPTR is

- Peer-reviewed
- bimonthly
- Rapid publication

Submit your manuscript at: editor@ajptr.com

